

USSN 09/890,335

Supplemental Amendment filed July 14, 2009


RECEIVED
CENTRAL FAX CENTER
JUL 14 2009Remarks/Arguments

No new matter is believed to be added by this Amendment.

The amendment to the chemical structure occurring in the paragraph spanning page 15 line 4 to page 16 line 2 is to correct a typographical error occurring within the phosphate group of the pictured chemical structure. Specifically, an oxygen atom is included with the phosphate group pictured as bound to group R_3 , so that " $P-R_3$ " is now " $P-O-R_3$ ". Support for this amendment is provided in the attached Appendix. While one skilled in the art may presume the presence of the oxygen atom in the structure as currently presented (see for instance a similar representation and typographical error in Mathew-van Holde's Biochemistry p. 303; Appendix), as Applicant is aware of the typographical error, Applicant corrects the error herein. This correction is also supported for instance by the disclosure at pages 15-16 that the chemical structure relates to the class of phospholipids having the formula defined therein, as phospholipids typically include the inserted oxygen. See for instance the Appendix included herein.

In the event that the Examiner has any questions or concerns regarding this application, the Examiner is invited to contact the below-signed representative by telephone to discuss.

Respectfully submitted,



Valerie Neymeyer-Tynkov
Reg. No. 46,956
Neymeyer-Tynkov LLC
20 N. Clark St. Suite 600
Chicago, IL 60602
Phone: 312/965-1969

PAGE 05/17
RECEIVED
CENTRAL FAX CENTER
JUL 14 2009

Appendix

Holum, Elements of General and Biological Chemistry: John Wiley & Sons, USA (1987) – pages 324, 325.

Stryer, Biochemistry: W.H. Freeman and Company, San Francisco, California (1981) – pages 208, 209.

Mathews and van Holde, Biochemistry: The Benjamin/Cummings Publishing Company, Redwood City, California (1990) – pages 303, 304.

SEVENTH EDITION

JOHN R. HOLM

ELEMENTS OF
GENERAL AND
BIOLOGICAL
CHEMISTRY

Production supervised by Linda R. Indig
Illustrations by John Balbalis with the assistance of
the Wiley Illustration Department
Design supervised by Ann Renzi
Photo researched by Stella Kupferberg
Manuscript edited by Vivian Danser under the
supervision of Deborah Herbert.
Cover photograph by Paul Silverman

Copyright © 1983, 1987, by John Wiley & Sons, Inc.

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of
this work beyond that permitted by Sections
107 and 108 of the 1976 United States Copyright
Act without the permission of the copyright
owner is unlawful. Requests for permission
or further information should be addressed to
the Permissions Department, John Wiley & Sons.

Library of Congress Cataloging in Publication Data:

Holum, John R.

Elements of general and biological chemistry.

Includes bibliographies and index.

1. Biological chemistry. I. Title.


QP514.2.H64 .1987 574.19'2 86-7827
ISBN 0-471-83331-4

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1


324 CHAPTER 15 LIPIDS

joined by a phosphate ester link to a small alcohol molecule. When this link is absent, the material is called phosphatidic acid.




Cephalin is from the Greek *kephale*, head. Cephalin is found in brain tissue.

The three principal phosphoglycerides are esters that are formed between phosphatidic acid and either choline, ethanolamine, or serine to give, respectively, phosphatidylcholine (lecithin), 2, phosphatidylethanolamine (cephalin), 3, and phosphatidylserine, 4.



As the structures of 2, 3, and 4 show, one part of each phosphoglyceride molecule is very polar because it carries full electrical charges. The remainder is nonpolar and hydrocarbon-like. These characteristics have important implications in understanding how phosphoglycerides are used to make cell membranes (Section 15.5).




Lecithin is from the Greek *lekithos*, egg yolk—a rich source of this phospholipid.

When pure, lecithin is a clear, waxy solid that is very hygroscopic. In air, it is quickly attacked by oxygen, which makes it turn brown in a few minutes. Lecithin is a powerful emulsifying agent for triacylglycerols, and this is why egg yolks, which contain it, are used to make the emulsions found in mayonnaise, ice cream, custards, and cake dough.


15.3 PHOSPHOLIPIDS 325

Plasmalogens. The plasmalogens make up another family of glycerol-based phospholipids, and they occur widely in the membranes of nerve cells and muscle cells. They differ from the other phosphoglycerides by the presence of an unsaturated ether group instead of an acyl group at one end of the glycerol unit.



Sphingolipids. The two types of sphingosine-based lipids or **sphingolipids** are the sphingomyelins and the cerebroside, and they are also important constituents of cell membranes. The sphingomyelins are phosphate diesters involving sphingosine. Their acyl units occur as acylamido parts, and they come from unusual fatty acids that are not found in neutral fats.

The cerebroside is not actually a phospholipid. Instead it is a **glycolipid**, a lipid with a sugar (i.e., glycosyl) unit and not a phosphate ester system. The sugar unit is usually that of D-galactose, or D-glucose, or amino derivatives of these.



Second Edition

BIOCHEMISTRY

STANFORD UNIVERSITY



W. H. FREEMAN AND COMPANY
San Francisco

DESIGNER: *Robert Ich*
ILLUSTRATOR: *Donna Salmon*
ILLUSTRATION COORDINATOR: *Andre W. Lorde*
PRODUCTION COORDINATOR: *William Murlack*
COMPOSITOR: *York Graphic Services*
PRINTER AND BINDER: *Arcata Book Group*

Library of Congress Cataloging in Publication Data

Stryer, Lubert.
Biochemistry.

Includes bibliographies and index.

1. Biological chemistry. I. Title. [DNLM:
1. Biochemistry. Q64 .S928b]
QP514.2.S65 1981 574.192 80-24699
ISBN 0-7167-1226-1

Copyright © 1975, 1981 by Lubert Stryer

No part of this book may be reproduced by any mechanical, photographic, or electronic process, or in the form of a phonographic recording, nor may it be stored in a retrieval system, transmitted, or otherwise copied for public or private use, without the written permission of the publisher.

Printed in the United States of America

9 8 7 6 5 4 3

208









Part I
CONFORMATION AND DYNAMICS

Figure 10-4

Absolute configuration of the glycerol 3-phosphate moiety of membrane lipids. (A) H and OH, attached to C-2, are in front of the plane of the page, whereas C-1 and C-3 are behind it. (B) Fischer representation of this structure. In a Fischer projection, horizontal bonds denote bonds in front, whereas vertical bonds denote bonds behind the plane of the page.



Now let us link some of these components to form phosphatidylcholine, a phosphoglyceride found in most membranes of higher organisms.



The structural formulas of the other principal phosphoglycerides—namely, phosphatidyl ethanolamine, phosphatidyl serine, phosphatidyl inositol, and diphosphatidyl glycerol—are given in Figure 10-5.

209

Chapter 10
INTRODUCTION TO MEMBRANES






Figure 10-5
Formulas of some phosphoglycerides.

Sphingomyelin is the only phospholipid in membranes that is not derived from glycerol. Instead, the backbone in sphingomyelin is *sphingosine*, an amino alcohol that contains a long, unsaturated hydrocarbon chain. In sphingomyelin, the amino group of the sphingosine backbone is linked to a fatty acid by an amide bond. In addition, the primary hydroxyl group of sphingosine is esterified to phosphoryl choline. As will be shown shortly, the conformation of sphingomyelin resembles that of phosphatidyl choline.



BIOCHEMISTRY

Christopher K. Mathews

Oregon State University

K. E. van Holde

Oregon State University

Illustration concepts by Audre W. Newman
with art contributions from Irving Geis

Cover

Dimer of *trp* repressor protein, with bound tryptophan, in blue. The protein binds to DNA and regulates expression of the *trp* genes that control tryptophan biosynthesis. Crystal structure by Paul Sigler et al.; image by Jane and David Richardson.

Frontispiece

Figure 11.15a The T state of aspartate transcarbamoylase, as determined by x-ray diffraction.

Sponsoring Editor: Diane Bowen
Developmental Editor: Susan Weisberg
Production Supervisor: Anne Friedman
Art Coordinator: Bruce Lundquist
Interior and Cover Designer: Gary Hlad
Diagram Artist: Wendy Calmenson
Copyeditor and Indexer: Mary Prescott
Principal Illustrator: Georg Klatt
Illustrators: Ken Miller, Elizabeth Morales-Denney, and Irene Imeldi
Proofreaders: Sheila Kennedy and Kathy Lee
Photo Research and Permissions: Rachel Menzi
Typesetter: York Graphic Services, Inc.
Film Preparation: Color Response, Inc.

Figures: 2.2, 2.3, 2.7, 2.9, 2.10, 2.11, 2.13, 2.14, 3.5, 4.2, 4.11, 4.12b, 4.14, 4.15, 4.22, 5.12, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.11, 6.12, 6.13a-c, 6.14, 6.17, 6.19, 6.21, 6.23a, b, and d, 6.25c and d, 7.2, 7.3b, 7.6, 7.9, 7.12, 7.13, 7.14, 7.15, 7.16, 7.17, 7.18, 7.21, 7.23 (H₂O molecule), 7.27, 7.28, 8.20, 8.21, 8.22, 8.23, 9.23, 15.7 copyright © Irving Geis.

Figure 4.15 Copyright © 1983 Dickerson and Geis.

Figure 10.13 Copyright © Stroud, Dickerson, and Geis.

Figure 11.15 Copyright © I. Geis and E. Gouaux, from coordinates by W. N. Lipscomb

Credits for photographs appear on pages xi-xii

Copyright © 1990 by The Benjamin Cummings Publishing Company, Inc.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written consent of the publisher. Printed in the United States of America. Published simultaneously in Canada.

Library of Congress Cataloging-in-Publication Data

Mathews, Christopher K., 1937—

Biochemistry / Christopher K. Mathews, K. E. van Holde; illustration concepts by Andre Newman with art contributions from Irving Geis.

p. cm.

Includes bibliographical references.

ISBN 0-8053-5015-2

I. Biochemistry. I. Van Holde, K. E. Kensal Edward, 1928—

II. Title.

DNLM: 1. Biochemistry. [QU 4 M4294b]

QP514.2.M384 1990

574.192—dc20

DNLM/DLC

for Library of Congress

89-17922

CLP

ISBN: 0-8053-5015-2

CDEFGHIJ-VII- 93210



The Benjamin Cummings Publishing Company
 390 Bridge Parkway
 Redwood City, CA 94065

The Lipid Constituents of Biological Membranes

All biological membranes contain lipids as major constituents. The molecules that play the dominant roles in membrane formation all have highly polar head groups and, in most cases, two hydrocarbon tails. There is a molecular sense to this: If a large head group is attached to a single hydrocarbon chain, the molecule is wedge shaped and will tend to form spherical micelles (Figure 9.5a). A double tail yields a roughly cylindrical molecule, which can easily pack in parallel to form extended sheets of membranes. As indicated in Figure 9.5b, such membranes will be bilayers, with the hydrophilic head groups facing outward into the aqueous regions on either side. A number of classes of membrane-forming lipids share this type of structure; they differ principally in the nature of the head group. We shall describe a few examples of each.

Glycerophospholipids

Glycerophospholipids (also called phosphoglycerides) are the major class of naturally occurring phospholipids, lipids with phosphate-containing head groups. These compounds make up a significant fraction of the membrane lipids throughout the bacterial, plant, and animal kingdoms. All can be considered to be derivatives of glycerol-3-phosphate. Carbon 2 in glycerol-3-phosphate is a chiral center, and the naturally occurring glycerophospholipids are derivatives of the L enantiomer. The general structure of this group of compounds is shown in Figure 9.6. In panel (a) is depicted the stereochemical configuration. Panel (b) shows the molecule in the manner we will generally use to represent membrane lipids, with the hydrophobic tails drawn to the right and the hydrophilic head group to the left. Usually, R_1 and R_2 are acyl side chains derived from the fatty acids; often one is saturated, the other unsaturated. The R_3 group varies greatly, and it is this that confers the greatest variation in properties among the glycerophos-




Figure 9.5
How molecular shape of lipids determines the structures they form. (a) The single tail of a fatty acid makes the molecule wedge-shaped, favoring micelle formation. (b) The multiple tails on membrane-forming lipids make the molecules more cylindrical, so that planar bilayer sheets can be formed.




Figure 9.6
Glycerophospholipid structure. (a) Stereochemical view of L-enantiomer. (b) A conventional visualization.

304




Figure 9.7

Examples of common glycerophospholipids. The hydrophobic R groups are indicated in yellow, the glycerol moiety in black, and the very hydrophilic head groups in blue. All may be considered derivatives of phosphatidic acid (a).